

Clinical and pathological effects of short-term cyanide repeated dosing to goats

B. Soto-Blanco,¹ B. L. Stegelmeier² and S. L. Górniak^{3,*}

¹ Departamento de Medicina Veterinária, Escola Superior de Agricultura de Mossoró, Mossoró, RN, Brazil

² USDA Agricultural Research Service, Poisonous Plant Research Laboratory, Logan, UT, USA

³ Centro de Pesquisas em Toxicologia Veterinária (CEPTOX), Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, SP, Brazil

Received 18 February 2004; Revised 7 July 2004; Accepted 19 July 2004

ABSTRACT: The purpose of this work is to determine and describe the effects of subacute cyanide toxicity to goats. Eight female goats were divided into two groups. The first group of five animals was treated with 8.0 mg KCN kg⁻¹ body weight day⁻¹ for seven consecutive days. The second group of three animals was treated with water as controls. Complete physical examination, including observation for behavior changes, was conducted before and after dosing. One treated animal was euthanized immediately after dosing. Later, two of the remaining treated animals and a control goat were euthanized after a 30-day recovery period. Euthanized animals were necropsied and tissues were collected and prepared for histologic studies. Clinical signs in treated goats were transient and included depression and lethargy, mild hyperpnea and hyperthermia, arrhythmias, abundant salivation, vocalizations, expiratory dyspnea, jerky movements and head pressing. Two goats developed convulsions after day 3 of treatment. One animal developed more permanent behavioral changes as she became less dominant and aggressive. Histologic changes included mild hepatocellular vacuolation and degeneration, mild vacuolation and swelling of the proximal convoluted tubules of the kidneys and spongiosis of the white matter (status spongiosis) of the cerebral white tracts, internal capsule, cerebellar peduncles, spinal cord and peripheral nerves. In summary, sub-lethal cyanide intoxication in goats resulted in behavioral changes, and during the treatment period animals showed delayed signs of toxicity. Significant histologic lesions in goats were observed and need to be characterized further. Copyright © 2005 John Wiley & Sons, Ltd.

KEY WORDS: behavioral sequelae; central nervous system; cyanide; cyanogenic plants; goats; myelin edema; status spongiosis

Introduction

Cyanogenic plants, such as cassava, sorghum and *Cynodon* grasses, have worldwide distribution and they are often ingested by animals (Tokarnia *et al.*, 2000). Many poisonings are caused by consumptions of relatively high doses consumed rapidly. These poisonings have been described extensively and it has been suggested that toxicity occurs through direct inhibition of the enzyme cytochrome oxidase, resulting in cellular hypoxia and fatal cytotoxic anoxia (Ballantyne, 1987). Aware of the acute effects of cyanide poisoning, farmers have adopted management strategies to reduce such toxicity. However, the effects of such sub-acute and chronic cyanide toxicity are unknown. Association of many different human diseases with chronic cyanide exposure suggests that

cyanide may cause many, yet unrecognized, problems that are of great importance for livestock kept on pastures containing cyanogenic plants.

Prolonged cyanide exposure has been associated with neurological and thyroidal disturbances in both humans and animals. Chronic cyanide exposure resulted in reduced body weight and weight gains in animals (Tewe *et al.*, 1984; Ibebunjo *et al.*, 1992; Soto-Blanco *et al.*, 2001a). Cyanide was found to disturb thyroid metabolism in rats (Philbrick *et al.*, 1979), rabbits (Ratnakumar and Rajan, 1992), pigs (Tewe *et al.*, 1984), goats (Bahri, 1987; Soto-Blanco *et al.*, 2001a) and humans (Adewusi and Akindahunsi, 1994). Studies in rats (Soto-Blanco *et al.*, 2002a) and goats (Soto-Blanco *et al.*, 2002b) chronically treated with cyanide have also described central nervous system (CNS) lesions.

The purpose of this work is to describe the effects of short-term toxicity to goats and compare them with those described in other species. Goats were chosen as the experimental species because several works evaluating cyanide toxicity in goats have been conducted in our laboratory (Soto-Blanco *et al.*, 2001a,b, 2002b; Soto-Blanco and Górniak, 2003, 2004), including their toxicokinetics (Sousa *et al.*, 2003).

* Correspondence to: S. L. Górniak, Centro de Pesquisas em Toxicologia Veterinária (CEPTOX), Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, Av. Prof. Dr. Orlando Marques de Paiva 87, 05508-000, São Paulo, SP, Brazil.

E-mail: gorniak@usp.br

Contract/grant sponsor: Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP); Contract/grant number: 99-08575-3.

Materials and Methods

Animals and Experimental Design

Eight, mixed-breed, mature female goats weighing 32–50 kg were fed 100 g of concentrate and sugar cane (*Saccharum officinarum* L.) *ad libitum*. The goats were divided into two groups. The first group (five animals) was treated with 8.0 mg of KCN (Merck, Darmstadt, Germany) kg body wt⁻¹ day⁻¹ diluted in tap water, divided into two daily doses, by gavage for seven consecutive days. The controls received only water for the same period.

Physical examinations and behavioral assessments were conducted before and after dosing. The term 'convulsions' was used only when severe muscular contractions occurred with loss of consciousness.

Cyanide Antidote

When treated goats became severely intoxicated (animals developed severe trembling and mild convulsion), sodium nitrite (10 g 100 ml⁻¹ of distilled water) at 20 mg kg body wt⁻¹ and sodium thiosulfate (20 g 100 ml⁻¹) at 500 mg kg⁻¹ were given i.v. as an antidote. The sodium thiosulfate treatment was repeated after 5 min.

Blood Biochemical Analysis

Blood samples were taken at the start of the study and at the end of the dosing periods and then at days 9, 10, 11 and 12. The blood samples were obtained from the jugular vein of each goat stored at -10 °C until analysis. Commercial kits were used for determination of glucose, cholesterol (Laborlab, Brazil), alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase, lactate dehydrogenase (LDH), gamma-glutamyltransferase (GGT), total protein, creatinine (Merck, Darmstadt) and urea nitrogen (Labtest, Brazil).

Plasma thiocyanate concentrations were determined by spectrophotometry following the method of Pettigrew and Fell (1972) but with minor modifications as described earlier (Soto-Blanco and Górniak, 2003).

Pathological Study

One treated goat (no. 1) was euthanized and necropsied immediately after the final dose. Two additional treated goats (nos 2 and 3) were euthanized and necropsied 30 days later. At necropsy the entire CNS, liver, spleen, kidneys, lungs, myocardium, pancreas, adrenal and thyroid glands were collected and fixed in Carnoy's solution. Paraffin-embedded sections were stained with H&E; CNS

sections were also stained with periodic acid-Schiff (PAS)/Luxol fast blue, Bodian silver and Holzers glial fiber stain.

Statistical Analysis

Data are reported as means ± SEM and were compared using a mixed model (PROC MIXED) for a repeated-measures design (SAS Statistical Software V8, 2000; SAS Institute, Cary, NC). The level of significance was set at $P < 0.05$.

Results

Clinical signs were transient and lasted up to 1 h after KCN administration. No changes were seen in control animals. After the initial dose, the treated goats were moved slowly or were reluctant to move, with hyperpnea and increased body temperature (up to 0.5 °C). Transient tachycardia also was found in most instances but other animals were normal or had mild bradycardia.

Other clinical signs that had variable presentation included abundant salivation, vocalizations, expiratory stridor, intention tremors, jerky head movement (usually quick upwards movement of the head and neck) and head pressing.

One goat (no. 2) developed convulsions immediately after the first daily KCN dose on day 3. Antidote was given and the convulsions subsided. Similar convulsions developed after treatment on subsequent days, so the second half of this animal's dose was not administered. However, the convulsions continued to occur. Another treated goat (no. 1) developed similar convulsions after the second dose of day 3 and was also treated with antidote. The same animal developed severe muscular contractions but not convulsions after the first dose on day 6. However, it quickly recovered and the antidote treatment was not necessary. The clinical effects were variable because treated animals responded differently. One very aggressive goat that tended to hit and bully the other goats gradually became less aggressive and this behavioral change was maintained through several weeks post-treatment.

The serum biochemical results are presented in Table 1. No significant inter-group differences were detected in serum glucose, creatinine, urea or cholesterol concentrations or in the activities of AST, ALT, GGT, or alkaline phosphatase. Plasma thiocyanate concentrations (Table 2) were higher in treated goats than in controls at all evaluated days but day 0, and the highest levels were found at day 9. Thiocyanate concentrations were lower in goat no. 2 than in other treated goats because this goat only received a single cyanide dose because of its high sensitivity to cyanide.

Table 1. Biochemical panel of female goats that received zero (control group) or 8.0 mg KCN kg⁻¹ day⁻¹ (treated group)

Group	Serum components ^a							
	GLU	CHOL	UN	CREA	AST	ALT	γ -GT	AP
Day 1								
Control	54.0 \pm 1.53	90.4 \pm 5.96	34.5 \pm 5.25	0.87 \pm 0.02	37.7 \pm 1.43	19.0 \pm 1.49	21.2 \pm 1.35	83.7 \pm 5.53
Treated	52.4 \pm 1.87	92.6 \pm 9.86	22.6 \pm 2.49	0.91 \pm 0.03	36.9 \pm 2.01	20.4 \pm 1.95	21.8 \pm 1.47	83.3 \pm 14.0
Day 7								
Control	48.3 \pm 1.86	84.8 \pm 8.59	35.1 \pm 2.91	0.91 \pm 0.02	36.7 \pm 0.54	20.0 \pm 2.35	20.6 \pm 1.35	75.2 \pm 4.84
Treated	50.2 \pm 2.92	91.8 \pm 10.4	32.6 \pm 0.90	0.91 \pm 0.06	36.2 \pm 1.59	23.0 \pm 1.29	22.6 \pm 0.92	61.3 \pm 10.8
Day 9								
Control	52.8 \pm 1.50	89.1 \pm 19.3	23.3 \pm 3.63	0.91 \pm 0.02	36.4 \pm 2.43	17.0 \pm 0.81	20.8 \pm 1.16	78.9 \pm 2.06
Treated	50.3 \pm 2.09	87.3 \pm 14.0	22.2 \pm 1.00	1.07 \pm 0.12	36.8 \pm 1.02	19.0 \pm 0.78	22.3 \pm 1.47	61.2 \pm 13.3
Day 30								
Control	52.1 \pm 1.85	90.0 \pm 16.7	41.2 \pm 2.95	0.94 \pm 0.01	42.1 \pm 0	17.0 \pm 0.81	23.4 \pm 0.29	67.8 \pm 4.13
Treated	45.7 \pm 3.49	93.6 \pm 9.10	34.7 \pm 3.11	0.97 \pm 0.08	42.5 \pm 0.77	21.4 \pm 1.79	23.3 \pm 0.96	59.5 \pm 6.19

^a Values are means \pm SEM. GLU, glucose (mg dl⁻¹); CHOL, cholesterol (mg dl⁻¹); UN, urea nitrogen (mg dl⁻¹); CREA, creatinine (mg dl⁻¹); AST, aspartate aminotransferase (U l⁻¹); ALT, alanine aminotransferase (U l⁻¹); γ -GT, γ -glutamyltransferase (U l⁻¹); AP, alkaline phosphatase (U l⁻¹).

Table 2. Plasma thiocyanate levels of female goats that received zero (control group) or 8.0 mg KCN kg⁻¹ day⁻¹ (treated group)



Period	Control ^a	Treated ^a	Goat no. 2
Day 0	7.56 \pm 0.34	7.04 \pm 0.40	7.56
Day 7	7.56 \pm 0.34	125.6 \pm 3.89*	79.4
Day 9			
Morning	8.40 \pm 0.24	145.9 \pm 31.6*	17.2
Afternoon	7.77 \pm 0.17	103.5 \pm 20.8*	14.7
Day 10			
Morning	8.61 \pm 0.26	28.0 \pm 9.55*	9.24
Afternoon	7.14 \pm 0.21	21.3 \pm 6.49*	9.24
Day 11			
Morning	8.19 \pm 0.17	15.1 \pm 6.00*	7.14
Afternoon	8.82 \pm 0.51	13.4 \pm 3.38*	8.34
Day 12	8.61 \pm 0.17	11.5 \pm 1.49*	8.08

^a Values are means \pm SEM; * $P < 0.05$, ANOVA followed by Duncan's test.

No significant gross lesions were found in any of the necropsied goats. Additionally, no microscopic changes were found in the control goat. In the treated animals, histologic changes included minimal hepatocellular vacuolation and swelling. This was most severe in periportal hepatocytes. The kidneys also had minimal vacuolation and swelling of the epithelial cells of the proximal convoluted tubules. The most striking lesion was that of mild spongiosis of the white matter of the cerebral white tracts, internal capsule and cerebellar peduncles (Figs 1 and 2). Occasionally, similar mild lesions were also present in the white tracts of the spinal cord and peripheral nerves.

Discussion

It has been suggested that the constant exposure to cyanide could increase the tolerance of the organism to its toxicity (D'Mello, 1987; Tokarnia *et al.*, 2000). On the other hand, the goats in this study did not present this

**Figure 1.** Cerebellar white matter showing spongiosis from a goat that received KCN for 7 days (H&E, 100 \times)**Figure 2.** Spongiosis on cerebellar white matter of a goat treated with KCN for 7 days (H&E, 1000 \times)

tolerance but they presented delayed signs of cyanide poisoning. Similar delayed signs were observed in goats treated with KCN for >120 days (Soto-Blanco *et al.*, 2001a), and in sheep that were treated with the

cyanogenic glucoside amygdalin (J. J. Villalba and F. D. Provenza, personal communication). This may be due to exhaustion of the mechanism of cyanide detoxification. A key step in cyanide detoxification is catalyzed by rhodanese because cyanide is bound irreversibly to sulfur from compounds such as thiosulphate and polythionates, producing thiocyanate. Although rhodanese is recycled and large amounts of cyanide can be metabolized, both cyanide and sulfite can inhibit rhodanese activity in the absence of thiosulfate (Bhatt and Linnell, 1987). Thus, the exposure to high, but not lethal, cyanide doses could deplete endogenous sulfur pools and partially impair cyanide detoxification. Another possibility is a neuronal accumulation of cyanide, which was observed earlier in *in vitro* studies (Borowitz *et al.*, 1994).

Long-term cyanogenic plant consumption by both men (Adewusi and Akindahunsi, 1994) and animals (Bahri, 1987; Kamalu and Agharanya, 1991) has been associated with the development of hypothyroidism. Thiocyanate, the main cyanide metabolite, is thought to be responsible because it competes with iodide in the thyroid gland (Delange and Ermans, 1996). Previous work with rats (Sousa *et al.*, 2002) and goats (Soto-Blanco *et al.*, 2001a) verified that the prolonged administration of KCN caused an increase in the number of resorption vacuoles in the follicles of the thyroid. However, no alterations were found in the thyroid glands of the treated animals in this study. This is likely to be due to the relatively short dosing period.

Tropical diabetes, or malnutrition-related diabetes mellitus, is a subclass of diabetes that has been associated with chronic cyanide exposure through consumption of cassava (McMillan and Geevarghese, 1979; Petersen, 2002). However, some researchers have not been able to reproduce cassava's diabetogenic effects using cyanide alone (Kamalu, 1991; Okolie and Osagie, 2000; Soto-Blanco *et al.*, 2001b). It may be that cassava contains a second diabetogenic toxin, or that this effect is a result of cyanide synergism with some contributing factor. Furthermore, epidemiological surveys in humans did not detect a relationship between cassava feeding and the development of pancreatic diabetes (Teuscher *et al.*, 1987; Vannasaeng *et al.*, 1988; Narendranathan and Cheriyan, 1994). This is probably a complex question and more studies are needed to determine fully whether cassava contains a yet unidentified diabetogenic toxin or is not involved in the etiology of tropical diabetes.

Nephrosis has been described also in rats (Ononogbu and Emole, 1978), dogs (Kamalu, 1993) and humans (Clark, 1936) fed cassava. Cassava treated dogs also developed periportal hepatocellular vacuolation (Kamalu, 1993). Recently Okolie and Osagie (1999) and Sousa *et al.* (2002) described similar degenerative changes in the liver and kidney from rabbits and rats treated with KCN. The goats in this study had minimal changes in both the liver and kidney that were not supported by the changes

in serum biochemical markers of liver and kidney disease. It is likely that they are not clinically significant at these doses and durations. Additional work is needed to determine if cyanide is nephro- and hepatotoxic to goats at higher doses or longer durations.

The main target system for cyanide's toxicity is inhibition of cellular respiration. The CNS is especially sensitive to many toxins that affect respiration. Neurons have relatively high metabolic rates with little capacity for anaerobic metabolism (D'Mello, 1987). Cyanide also may be directly neurotoxic. It has been shown to interfere with several neurotransmitters including γ -aminobutyric acid (GABA), glutamic acid (Cassel *et al.*, 1991), acetylcholine (Owasoya and Iramain, 1980), dopamine (Cassel *et al.*, 1995), excitatory amino acids and nitric oxide (Gunasekar *et al.*, 1996). Furthermore, Borowitz *et al.* (1997) suggested that cyanide also would be an endogenous neurotransmitter.

The change of behavior seen in one treated goat in this study was interesting, and to our knowledge no similar finding has been described in livestock. Some sequelae of cyanide toxicity in humans have been associated with damage to the basal ganglia, including Parkinsonian symptoms (Finelli, 1981; Uitti *et al.*, 1985; Carella *et al.*, 1988; Rosemberg *et al.*, 1989; Feldman and Feldman, 1990; Messing, 1991), phobic anxiety (Nicholson and Vincenti, 1994) and psychosis (El Ghawabi *et al.*, 1975; Kales *et al.*, 1997). Furthermore, cyanide intoxication has been linked to personality change in a man (Uitti *et al.*, 1985). In rodents, motor and cognitive function (D'Mello, 1987) and memory (Mathangi and Namasivayam, 2000) have been hypothesized as being affected by cyanide.

The vacuolation of neuropil (status spongiosis) seen in the myelin tracts in the cerebral and cerebellar white matter is suggestive of myelinic edema. Additional work, including electron microscopy, is needed to characterize better this lesion. In an earlier experiment, goats chronically treated with KCN developed axonal dystrophy (spheroids) and spongiosis and gliosis in the medulla oblongata (Figs 3 and 4), gliosis in the pons and Purkinje cell degeneration in the cerebellum (Soto-Blanco *et al.*, 2002b). The differences in pathological findings are likely to be due to differences in dose, duration and physiological status, because the earlier study used growing male goats that received lower doses of KCN (up to $3.0 \text{ mg kg}^{-1} \text{ day}^{-1}$) for 5 months (Soto-Blanco *et al.*, 2002b).

Status spongiosis is the term applied to the appearance of spongy nervous tissue on light microscopy (Adornato and Lampet, 1971). Various spongiform myelinopathies in ruminants (similar to found in our goats) were described as resulting from poisoning by hexachlorophane (Hall and Reid, 1974), including: *Stypandra glauca* (Whittington *et al.*, 1988), *S. imbricata* (Main *et al.*, 1981), *Diplodia maydis* (Prozesky *et al.*, 1994) and *Helichrysum argyrosphaerum* (Van der Lugt *et al.*, 1996). Additional



Figure 3. Medulla oblongata of a goat that received KCN for 7 days, showing severe spongiosis on white tracts (H&E, 100x)




Figure 4. Spongiosis on white tracts in the medulla oblongata of a goat treated with KCN for 7 days (H&E, 1000x)

comparative and mechanistic studies are needed for better characterization of these lesions, to determine how they develop and if the process is reversible.

In summary, we found that sub-lethal cyanide doses given repeatedly for 7 days cause significant clinical and neurologic lesions. Such cyanide exposure is certainly dangerous and is likely to compromise significantly the animal's health.

Acknowledgments—The authors wish to thank Breno Shumacher-Henrique for help with clinical examinations and veterinary care during treatment, Leonila E. R. Rasputini for biochemical determinations and Estevao Belloni, Marco Faustino and Adilson Baladore for animal care and handling. This work was part of B. S.'s PhD thesis of Benito Soto-Blanco at the Department of Pathology, School of Veterinary Medicine, University of São Paulo, Brazil.

References

Adewusi SRA, Akindahunsi AA. 1994. Cassava processing, consumption, and cyanide toxicity. *J. Toxicol. Environ. Health* **43**: 13–23.

- Adornato B, Lampert P. 1971. Status spongiosis of nervous tissue. *Acta Neuropathol. (Berlin)* **19**: 271–289.
- Bahri S. 1987. Chronic effect of cassava leaves on the plasma thiocyanate and on the performance of the goat thyroid gland. *Penyakit Hewan* **19**: 88–90.
- Ballantyne B. 1987. Toxicology of cyanides. In *Clinical and Experimental Toxicology of Cyanides*, Ballantyne B, Marrs TC (eds). Wright: Bristol; 41–126.
- Bhatt HR, Linnell JC. 1987. The role of rhodanese in cyanide detoxification: its possible use in acute cyanide poisoning in man. In *Clinical and Experimental Toxicology of Cyanides*, Ballantyne B, Marrs TC (eds). Wright: Bristol; 440–450.
- Borowitz JL, Rathinavelu A, Kanthasamy A, Wilsbacher J, Isom GE. 1994. Accumulation of labeled cyanide in neuronal tissue. *Toxicol. Appl. Pharmacol.* **129**: 80–85.
- Borowitz JL, Gunasekar PG, Isom GE. 1997. Hydrogen cyanide generation by μ -opioid receptor activation: possible neuromodulatory role of endogenous cyanide. *Brain Res.* **768**: 294–300.
- Carella F, Grassi MP, Savoirdo M, Contri P, Rapuzzi B, Mangoni A. 1988. Dystonic-Parkinsonian syndrome after cyanide poisoning: clinical and MRI findings. *J. Neurol. Neurosurg. Psychiatry* **51**: 1345–1348.
- Cassel G, Karlsson L, Sellstrom A. 1991. On the inhibition of glutamic acid decarboxylase and gamma-aminobutyric acid transaminase by sodium cyanide. *Pharmacol. Toxicol.* **69**: 238–241.
- Cassel G, Koch M, Tiger G. 1995. The effects of cyanide on the extracellular levels of dopamine, 3,4-dihydroxyphenylacetic acid, homovanillic acid, 5-hydroxyindoleacetic acid and inositol phospholipid breakdown in the brain. *Neurotoxicology* **16**: 73–82.
- Clark A. 1936. Report on effects of certain poisons contained in food plants of West Africa upon the health of the natives races. *J. Trop. Med. Hyg.* **39**: 269–276.
- Delange FM, Ermans AM. 1996. Iodine deficiency. In *Werner and Ingbar's The Thyroid: a fundamental and clinical text* (7th edn), Braverman LE, Utiger RD (eds). Lippincott-Raven: Philadelphia; 296–316.
- D'Mello GD. 1987. Neuropathological and behavioural sequelae of acute cyanide toxicosis in animal species. In *Clinical and Experimental Toxicology of Cyanides*, Ballantyne B, Marrs TC (eds). Wright: Bristol; 156–183.
- El Ghawabi SH, Gaafar MA, El-Saharti AA, Ahmed SH, Malash KK, Fares R. 1975. Chronic cyanide exposure: a clinical, radioisotope, and laboratory study. *Br. J. Ind Med.* **32**: 215–219.
- Feldman JM, Feldman MD. 1990. Sequelae of attempted suicide by cyanide ingestion: a case report. *Psychiatry Med.* **20**: 173–179.
- Finelli PF. 1981. Changes in the basal ganglia following cyanide poisoning. *J. Comput. Assist. Tomogr.* **5**: 755–756.
- Gunasekar PG, Sun PW, Kanthasamy AG, Borowitz JL, Isom GE. 1996. Cyanide-induced neurotoxicity involves nitric oxide and reactive oxygen species generation after *N*-methyl-D-aspartate receptor activation. *J. Pharmacol. Exp. Ther.* **277**: 150–155.
- Hall GA, Reid IM. 1974. The effects of hexachlorophane on the nervous system of sheep. *J. Pathol.* **114**: 241–246.
- Ibebunjo C, Kamalu BP, Ihemelandu EC. 1992. Comparison of the effects of cassava (*Manihot esculenta* Crantz), organic cyanide and inorganic cyanide on muscle and bone development in a Nigerian breed of dog. *Br. J. Nutr.* **68**: 483–491.
- Kales SN, Dinklage D, Dickey J, Goldman RH. 1997. Paranoid psychosis after exposure to cyanide. *Arch. Environ. Health* **52**: 245–246.
- Kamalu BP. 1991. The effect of a nutritionally-balanced cassava (*Manihot esculenta* Crantz) diet on endocrine function using the dog as a model. 1. Pancreas. *Br. J. Nutr.* **65**: 365–372.
- Kamalu BP. 1993. Pathological changes in growing dogs fed on a balanced cassava (*Manihot esculenta* Crantz). *Br. J. Nutr.* **69**: 921–934.
- Kamalu BP, Agharanya JC. 1991. The effect of a nutritionally-balanced cassava (*Manihot esculenta* Crantz) diet on endocrine function using the dog as a model. 2. Thyroid. *Br. J. Nutr.* **65**: 373–379.
- Main DC, Slatter DH, Huxtable CR, Constable IC, Dorling PR. 1981. *Stypandra imbricata* ('blindgrass') toxicosis in goats and sheep – clinical and pathologic findings in 4 field cases. *Aust. Vet. J.* **57**: 132–135.
- Mathangi DC, Namasivayam A. 2000. Effect of chronic cyanide intoxication on memory in albino rats. *Food Chem. Toxicol.* **38**: 51–55.

- McMillan DE, Geevarghese PJ. 1979. Dietary cyanide and tropical malnutrition diabetes. *Diabetes Care* **2**: 202–208.
- Messing B. 1991. Extrapyramidal disturbances after cyanide poisoning (first MRT-investigation of the brain). *J. Neural Transm. Suppl.* **33**: 141–147.
- Narendranathan M, Cheriyan A. 1994. Lack of association between cassava consumption and tropical pancreatitis syndrome. *J. Gastroenterol. Hepatol.* **9**: 282–285.
- Nicholson PJ, Vincenti GE. 1994. A case of phobic anxiety related to the inability to smell cyanide. *Occup. Med. Oxf.* **44**: 107–108.
- Okolie NP, Osagie AU. 1999. Liver and kidney lesions and associated enzyme changes induced in rabbits by chronic cyanide exposure. *Food Chem. Toxicol.* **37**: 745–750.
- Okolie NP, Osagie AU. 2000. Differential effects of chronic cyanide intoxication on heart, lung and pancreatic tissues. *Food Chem. Toxicol.* **38**: 543–548.
- Ononogbu IC, Emole I. 1978. The effect of garri on rat plasma cholesterol. *Atherosclerosis* **31**: 101–104.
- Owasoyo JO, Iramain CA. 1980. Acetylcholinesterase activity in rat brain: effect of acute cyanide intoxication. *Toxicol. Lett.* **6**: 1–3.
- Petersen JM. 2002. Tropical pancreatitis. *J. Clin. Gastroenterol.* **35**: 61–66.
- Pettigrew AR, Fell GS. 1972. Simplified colorimetric determination of thiocyanate in biological fluids and its application to investigation of the toxic amblyopias. *Clin. Chem.* **18**: 996–1000.
- Philbrick DJ, Hopkins JB, Hill DC, Alexander JC, Thomson RG. 1979. Effects of prolonged cyanide and thiocyanate feeding in rats. *J. Toxicol. Environ. Health* **5**: 579–592.
- Prozesky L, Kellerman TS, Swart DP, Maartens BP, Schults RA. 1994. Perinatal mortality in lambs of ewes exposed to cultures of *Diplodia maydis* (= *Stenocarpella maydis*) during gestation. A study of the central-nervous-system lesions. *Onderstepoort. J. Vet. Res.* **61**: 247–253.
- Ratnakumar JN, Rajan A. 1992. Goitrogenic effect of cassava in broiler rabbits. *Ind. J. Animal Sci.* **62**: 670–676.
- Rosemberg NL, Myers JA, Martin WRW. 1989. Cyanide-induced Parkinsonism: clinical, mark, and 6-fluorodopa PET studies. *Neurology* **39**: 142–144.
- Soto-Blanco B, Gorniak SL. 2003. Milk transfer of cyanide and thiocyanate: cyanide exposure by lactation in goats. *Vet. Res.* **34**: 213–220.
- Soto-Blanco B, Gorniak SL. 2004. Prenatal toxicity of cyanide in goats – a model for teratological studies in ruminants. In *Poisonous Plants and Related Toxins*, Acamovic T, Stewart CS, Pennycott TW (eds). CABI Publishing: Oxon; 425–430.
- Soto-Blanco B, Gorniak SL, Kimura ET. 2001a. Physiopathological effects of chronic cyanide administration to growing goats – a model for cyanogenic plants ingestion. *Vet. Res. Common* **25**: 379–389.
- Soto-Blanco B, Sousa AB, Manzano H, Guerra JL, Gorniak SL. 2001b. Does prolonged cyanide exposure have a diabetogenic effect? *Vet. Human Toxicol.* **43**: 106–108.
- Soto-Blanco B, Maiorka PC, Gorniak SL. 2002a. Effects of long-term low-dose cyanide administration to rats. *Ecotoxicol. Environ. Safety* **53**: 37–41.
- Soto-Blanco B, Maiorka PC, Gorniak SL. 2002b. Neuropathologic study of long-term cyanide administration to goats. *Food Chem. Toxicol.* **40**: 1693–1698.
- Sousa AB, Soto-Blanco B, Guerra JL, Kimura ET, Gorniak SL. 2002. Does prolonged oral exposure to cyanide promote hepatotoxicity and nephrotoxicity? *Toxicology* **174**: 87–95.
- Sousa AB, Manzano H, Soto-Blanco B, Gorniak SL. 2003. Toxicokinetics of cyanide in rats, pigs and goats after oral dosing with potassium cyanide. *Arch. Toxicol.* **77**: 330–334.
- Teuscher T, Baillod P, Rosman JB, Teuscher A. 1987. Absence of diabetes in a rural West African population with a high carbohydrate/cassava diet. *Lancet* **1**: 765–768.
- Tewe OO, Afolabi AO, Grissom FEG, Littleton GK, Oke OL. 1984. Effect of varying dietary cyanide levels on serum thyroxine and protein metabolites in pigs. *Nutr. Rep. Int.* **30**: 1249–1253.
- Tokarnia CH, Döbereiner J, Peixoto PV. 2000. *Plantas Tóxicas do Brasil*. Helianthus: Rio de Janeiro; 215–221.
- Uitti RJ, Rajput AH, Ashenhurst EM, Rozdilsky B. 1985. Cyanide-induced parkinsonism: a clinicopathologic report. *Neurology* **35**: 921–925.
- Van der Lugt JJ, Oliver J, Jordaan P. 1996. Status spongiosis, optic neuropathy, and retinal degeneration in *Helichrysum argyrosphaerum* poisoning in sheep and a goat. *Vet. Pathol.* **33**: 495–502.
- Vannasaeng S, Nitiyanant W, Vichayanrat A. 1988. Case-control study on risk factors associated with fibrocalculous pancreatic diabetes. *Diabetes Med.* **5**: 835–839.
- Whittington RJ, Searson JE, Whittaker SJ, Glastonbury JRW. 1988. Blindness in goats following ingestion of *Stypandra glauca*. *Aust. Vet. J.* **65**: 176–181.